

RYEQO® INFORMATION ABOUT YOUR ENDOMETRIOSIS TREATMENT



This leaflet was created by Gedeon Richter and is ONLY for patients who have been prescribed Ryeqo®.

This material should not be used as a substitute for the Consumer Medicine Information (CMI) for Ryeqo® or professional medical advice. Please refer to the CMI for detailed information on this medicine.

WHY HAVE I REEN PRESCRIBED RYEQO?

You have been prescribed Ryego to help treat your symptoms of endometriosis. The information in this booklet is here to guide and support you through your new treatment journey.



FACT

You are not alone - 1 in 7 women and those assigned female at birth are estimated to have endometriosis in Australia.2


WHAT IS ENDOMETRIOSIS?

Endometriosis is a condition where the inner lining of your uterus grows outside your uterus.^{3,4} Hormonal changes cause this tissue to thicken, break down and bleed during each menstrual cycle causing inflammation. Over time, the broken down tissue may become trapped in your pelvis causing scarring and adhesions.⁵

Common symptoms of endometriosis include chronic pelvic pain and/or struggling to conceive. The pain associated with endometriosis often occurs one or two days before your period and lasts until the end of your period.^{3,5} Other symptoms include pain during sex, painful bowel movements and gastrointestinal complaints.5,6

WHAT IS RYEQO?1

Ryeqo is a treatment for women who have already had other medical or surgical treatment for their endometriosis. Ryeqo contains relugolix, and the hormones, estradiol and norethisterone acetate. Relugolix reduces the amount of estrogen and progesterone circulating in the body. Estradiol and norethisterone acetate are combined with Relugolix to relieve your symptoms of endometriosis while helping to protect the strength of your bones.




WILL RYEQO CURE MY ENDOMETRIOSIS?

There is currently no cure for endometriosis, but there are a number of treatments available to manage the symptoms of endometriosis.^{3,6} Ryeqo is a non-surgical option that helps reduce the impact symptoms have on your life by offering effective relief from your pain.1

It has been shown that treatment with Ryego leads to substantial and sustained improvements in the quality of life for women with endometriosis. This is achieved by Ryego reducing their symptoms and the distress they cause.⁷

This includes not only reducing the experienced pain, but also improving energy and mood, the ability to carry out physical and social activities, and sexual function – all of which lead to a better sense of wellbeing.^{1,7}

Whilst Ryego effectively reduces the symptoms of endometriosis for the majority of women+, it is not effective for everyone, and some women+ will experience side effects. Your doctor or nurse can provide support if Ryego doesn't offer enough symptom relief from endometriosis, or if you experience any side effects. Alternatively, please contact Gedeon Richter for support using the contact details on the back page of this booklet.



RYEQO AND BONE MINERAL DENSITY (BMD)

- BMD is a measure of the amount of minerals contained in your bones.
 Measuring BMD using an X-ray test called a DXA scan can indicate the strength and health of your bones.
- While the estradiol and norethisterone acetate component of Ryeqo is designed to mitigate the risk of BMD loss, if you have existing bone health issues, such as osteoporosis, you should not take Ryeqo.⁸
- It is therefore recommended that your doctor check that your bones are healthy before starting Ryeqo, and monitor your bone health every year that you take Ryeqo with an annual DXA scan.¹



Your doctor is recommended to order a DXA scan to check your BMD when you start Ryeqo and then monitor it with a DXA scan each year after that.¹

HOW DO I TAKE RYEQO?8

It's simple. Ryeqo is a small tablet that only needs to be taken **once** a day.



IT IS RECOMMENDED THAT YOU:



Start taking Ryeqo within the first 5 days after the start of bleeding due to your period.




Keep taking it daily at about the same time of day



WHAT SHOULD I EXPECT FROM TREATMENT?

With Ryego, most women see a significant effect on their endometriosis symptoms, as soon as four weeks. If it takes a little longer for you to see improvements, this is completely normal.

Ryego works on your hormone levels and this may disrupt your usual menstrual cycle. In most cases, Ryego will reduce menstrual bleeding within the first 2 months of treatment, and many patients may eventually experience no menstrual bleeding at all.1



WILL I EXPERIENCE **ANY SIDE EFFECTS?8**

As with any treatment, side effects are a possibility – however, when Ryego patients were compared to those who did not receive any medical treatment for endometriosis, there were no significant side effects experienced.

The most common side effects include headache and hot flushes. Less common side effects include: dizziness, nausea, excessive, irregular, or prolonged bleeding from the womb (uterine bleeding), hair loss, decreased interest in sex, irritability, increased sweating, a lump in the breast tissue (breast cyst), indigestion. You should talk to your doctor if you experience any side effects while taking Ryego.

WHAT ABOUT RYEQO AND CONTRACEPTION?8

Firstly, you should not take Ryeqo if you are pregnant or breastfeeding. You will also need to stop taking other hormonal treatments, including hormonal contraception, before starting Ryeqo.

After 1 month of continuous use, Ryeqo will have a contraceptive effect (will prevent pregnancy). You are therefore unlikely to become pregnant while you take Ryeqo as recommended.

However, should you want to become pregnant in the near future, don't worry: your fertility will return rapidly after you stop taking Ryeqo.



Non-hormonal birth control (e.g. condoms) should be used during the first month after starting Ryeqo.⁸

WHAT IF I MISS A TABLET?8

If you miss a tablet, don't worry, just take the missed tablet as soon as you remember, you should then resume taking your tablet the next day as usual. Do not take a double dose to make up for a missed tablet. If you miss doses for 2 or more days, use a non-hormonal method of contraception (such as condoms) for the next 7 days.

WHERE CAN I GET FURTHER INFORMATION?

If you have any concerns or questions about endometriosis or Ryego, please speak to your healthcare professional.



For detailed information:

Please refer to the Ryeqo° Consumer Medicine Information for detailed information on using this medication. The CMI is available from the Therapeutic Goods Association using this QR code.

Alternatively, please contact us:

Phone 1300 GEDEON (1300 433 033)

e-mail medinfo@gedeonrichter.com.au



References

- 1. Ryego Approved Product Information. January 2024.
- 2. Australian Institute of Health and Welfare (2023) Endometriosis, AIHW, Australian Government, accessed 22 January 2024.
- **3.** Office on Women's Health in the U.S. Department of Health and Human Services. 2021. Endometriosis. Accessed November 2023. Available from: https://www.womenshealth.gov/a-z-topics/endometriosis.
- **4.** Wolters Kluwer N.V. UpToDate, Inc. 2023. Patient education: Endometriosis (Beyond the Basics). Accessed November 2023. Available from: https://www.uptodate.com/contents/endometriosis-beyond-the-basics#H3.
- **5.** Healthline Media LLC. 2023. Endometriosis. Accessed November 2023. Available from: https://www.healthline.com/health/endometriosis?
- 6. NHS.UK. 2022. Endometriosis-Treatment. Accessed November 2023. Available from:

https://www.nhs.uk/conditions/endometriosis/treatment/.

- 7. Giudice LC et al. Lancet. 2022;399:2267-2279.
- 8. Ryeqo Consumer Medicine Information. January 2024.

Tell your doctor as soon as possible if you experience any side effects using Ryeqo*.

Ryeqo* is a registered trademark of Gedeon Richter. Gedeon Richter Australia Pty Ltd ABN 98 602 550 274. Suite 902/15 Blue St, North Sydney NSW 2060. Date of approval: September 2024. COMS-0082-FEB-24.